Appendix 1. Secondary outcomes measured as part of this RCT.

Secondary Outcome	Definition
Incremental cost	Incremental cost, defined as the cost associated with respective
	video interventions (development, start-up, implementation, condom
	use, intervention – see Table 2 for details) per individual who
	reported no sex or sex with a condom during the follow-up period.
Female condomless	Frequency of men, defined as number of men who reported
sex	condomless vaginal or anal sex with a woman divided by the total
	number of men who viewed the video in that arm.
Male condomless sex	Frequency of men, defined as number of men who reported
	condomless anal sex with a man divided by the total number of men
	who viewed the video in that arm
Post-video condomless	Frequency of men, defined as number of men who reported
sex	condomless vaginal or anal sex with any partner immediately
	following the video intervention divided by the total number of men
	who viewed the video in that arm
Frequency of sex acts	Frequency of men, defined as the number of men who had decreased
	total number of sex acts in the three weeks following the
	intervention compared to the three weeks immediately preceding the
	intervention in that arm
Condom use social	Frequency of men, defined as number of men who report higher
norms	levels of social norms when comparing their pre-intervention and
	post-intervention condom use norms*
Condom self-efficacy	Frequency of men, defined as number of men who had an increase
	in self-efficacy when comparing their pre-intervention and post-
	intervention self-efficacy**
Condom negotiation	Frequency of men, defined as the number of men who attempted to
	convince an unwilling partner to use a condom immediately
	following the video intervention divided by the total number of men
	who viewed the video in that arm
HIV testing	Frequency of men, defined as the number of men who reported
	being tested for HIV during the interval between watching the video
	and following up compared to the number of men who followed up
STI testing	Frequency of men, defined as the number of men who reported
	being tested for STIs (excluding HIV) during the interval between
	watching the video and following up compared to the number of
	men who followed up

<sup>\*</sup>Condom use social norms will be measured using six survey items that are each on a five point Likert scale. Increased condom use social norms will be defined as having an increase from baseline in any two of these six survey items and dichotomized accordingly. The condom use social norm outcome will be assessed in the entire group as well as the subgroup of men who were referred by their friends.

<sup>\*\*</sup>Self-efficacy will be measured using seven survey items that are each on a five point Likert scale. Increased self-efficacy will be defined as having an increase from baseline in any two of

these seven survey items and dichotomized accordingly. The self-efficacy outcome will be assessed in the entire group as well as the subgroup of men who were referred by their friends.